Surface Inversions, Atmospheric Stability, and Spray Drift

Surface Temperature Inversions:

- Are very common
- Are easy to recognize
- Affect the dispersal of very small spray droplets suspended in the air
- Do not increase the amount of off-site movement
- Can increase the potential for offsite affects & the distance at which affects can be observed

Atmospheric Stability

- Inversions cause STABLE atmospheric conditions
- The concept of atmospheric stability helps understand how inversions affect drift
- Close to the ground, atmospheric stability changes regularly between STABLE, NEUTRAL, & UNSTABLE

The large scale trend in the atmosphere is that temperature decreases

70° F with height


75° F


80° F


Where there is a decrease in temperature with height greater than the adiabatic rate the atmosphere is UNSTABLE

75°F

80°F







A temperature inversion is an

60 °F area where temperature
increases with height.

55 °F

50 °F

It can begin at the ground.

45 °F

40 °F

50°F
48°F
46°F
44°F
42°F
40°F
38°F
These aloft inversions will be discussed at the end of the program
42°F
44°F
44°F
44°F
44°F
46°F
48°F


When temperature increases


with height the atmosphere is
STABLE.


55°F


Vertical mixing of the air and
dispersion of small droplets is
suppressed.


40°F



















































How to recognize a surface inversion

- Sunset to just after sunrise
- Windless to low wind conditions (<2-3 mph)
- Clear to partly cloudy skies
- Ground fog (if sufficient humidity exists)
- Dust hanging over a roadway
- Smoke from a chimney forming a layer
- Dew or frost (if sufficient humidity exists)





How to recognize the <u>potential</u> for a surface inversion

- Sun is getting low in the sky
- Wind is becoming light and variable
- Clear to partly cloudy skies

Surface Inversions don't:


• Increase the amount of spray droplets moving off-site


Surface Inversions can:


- Decrease the dispersion of droplets too small to quickly settle out.
- Result in a higher air concentration of these small droplets.
- Increase the potential for off-target effects.
- Increase the distance at which off-target effects can be observed.
- Increase the size of the area affected.
- Cause the direction of drift to be unpredictable


Reducing the effects of Surface Inversions on Spray Drift

- Minimizing production of very small drops
- Using equipment the minimizes the number of small drops suspended in the air
- Morning applications are likely to have shorter exposure to STABLE conditions than evening applications









Neutral conditions can be the best time to spray

- Wind direction if often consistent
- Good dispersion of droplets too small to quickly settle out.

50°F **Aloft Inversions** 48°F 44°F • Are not likely to effect applications to 42°F agricultural crops 40°F • Low altitude aloft inversions can affect 38°F aerial forestry application 40°F • Low altitude aloft inversions are often surface inversions in the process of 44°F dissipating & are therefore short lived 46°F



Surface Temperature Inversions:

- Are very common
- Are easy to recognize
- Affect the dispersal of very small spray droplets suspended in the air
- Do not increase the amount of off-site movement
- Can increase the potential for offsite affects & the distance at which affects can be observed